

## Exercise 3 – National dataset development by adjustment

With the information obtained in the training, how would you create datasets that are representative for Sri Lanka for one of the following options, Rice production and milling, Tea production and processing or Dairy production?

For a good overview of existing datasets, you can consult <https://nexus.openlca.org>.

Below you can find examples of sources for **Rice datasets**.

[https://www.lcacommons.gov/discovery/search?loc=&max=50&year=&dtype=&hfacet=%22Agriculture%2C+forestry+and+fishing%7EISIC+0112%3A+Growing+of+rice%22&SYNCHRONIZER\\_URI=%2Fdiscovery%2Fsearch&qlookup=&hfacetCat=&equip=&provider=&id=&crop=&SYNCHRONIZER\\_TOKEN=6d509f8d-d1d5-4f62-93e6-c141e6984ed0](https://www.lcacommons.gov/discovery/search?loc=&max=50&year=&dtype=&hfacet=%22Agriculture%2C+forestry+and+fishing%7EISIC+0112%3A+Growing+of+rice%22&SYNCHRONIZER_URI=%2Fdiscovery%2Fsearch&qlookup=&hfacetCat=&equip=&provider=&id=&crop=&SYNCHRONIZER_TOKEN=6d509f8d-d1d5-4f62-93e6-c141e6984ed0)

The screenshot displays the USDA National Agricultural Library search interface. At the top, the USDA logo and 'National Agricultural Library' are visible, along with the text 'United States Department of Agriculture'. A navigation bar includes links for Home, Data & Tools, Data Documentation, About, Contact Us, and Help. Below the navigation bar, there are buttons for 'Submit Data' and 'Downloads'. A search bar is present with the text 'Search for (Previous Searches) (New search)' and a placeholder 'For example: corn grain'. The search results section shows '6 records found (Showing items 1 through 6)'. The results list includes: 'rice; at harvest, at farm, wet basis US-CA', 'rice; at harvest, at farm, wet basis US-AR', 'rice; at harvest, at farm, wet basis US-LA', 'rice; at harvest, at farm, wet basis US-MS', 'rice; at harvest, at farm, wet basis US-TX', and 'rice; at harvest, at farm, wet basis US-MO'. On the left side, a 'Product Category' filter is shown with a tree structure. The 'Agriculture, forestry and fishing' category is expanded, showing sub-categories like 'ISIC 0111: Growing of cereals (except rice), leguminous crops and oil seeds' (93), 'ISIC 0112: Growing of rice' (6), 'ISIC 0116: Growing of fibre crops' (12), 'ISIC 0145: Raising of swine/pigs' (19), and 'ISIC 0161: Support activities for crop production' (31946). Other categories include 'Construction' (9).



Sri Lanka and IRRI started collaborating in 1960 through training and exchange of rice varieties. In 1967, an agreement between Sri Lanka and the [Ford Foundation](#) led to a two-year program between IRRI and the country's Department of Agriculture where its scientists eventually trained at IRRI. Renewed in 1969, the program also included technology transfer activities.

Different projects were conducted with the collaborative efforts of IRRI and partner institutions in Sri Lanka, one of which is a collaboration with the country's Department of Agriculture (DOASL) and the [United States Agency for International Aid \(USAID\)](#). They undertook a Rice Research project where Sri Lankans were provided with nearly 25 graduate scholarships to different universities in the US, UK, and the Philippines, as well as 100 short-term trainings in rice production, and cropping systems. Moreover, the project also provided research equipment and infrastructure development in rice research stations, and included as a component the improvement of rice varieties.

In 2007, DOASL and IRRI held a collaborative workplan meeting where they identified some areas for mutual cooperation. This included improving existing rice varieties in terms of quality and yield, enhancing conservation of rice genetic resources, increasing labor productivity, and strengthening the delivery and impact of technology through good extension models. Similar with past agreements, DOASL continued to serve as the clearing house for IRRI's activities in Sri Lanka. Both institutions continue to find ways and means of improving collaboration including funding support.



## 1.1.1 RICE CULTIVATION

### RICE

Rice is the single most important crop occupying 34 percent (0.77 /million ha) of the total cultivated area in Sri Lanka. On average 560,000 ha are cultivated during maha and 310,000 ha during yala making the average annual extent sown with rice to about 870,000 ha. About 1.8 million farm families are engaged in paddy cultivation island-wide. Sri Lanka currently produces 2.7 million t of rough rice annually and satisfies around 95 percent of the domestic requirement. Rice provides 45% total calorie and 40% total protein requirement of an average Sri Lankan. The per capita consumption of rice fluctuates around 100 kg per year depending on the price of rice, bread and wheat flour.

The current cost of production of rough rice is Rs. 8.57 per kg. The cost of labor, farm power and tradable inputs constitutes 55%, 23% and 23% respectively.

Field water requirement for a rice crop depends mainly on the growth duration of the crop and its growing environment. It is calculated that about 30-40% of the total water supplied to an irrigated crop is often supplied before the establishment of the rice crop and the amount is dependent on the soil drainage class, weed density and time taken for land preparation. Time taken for land preparation could be minimised to about 2 weeks using total killing herbicides (e.g. Paraquat) which also would help to reduce one tillage operation and conserve irrigation water.

#### Recommended Herbicides for Rice Weeds

##### One shot herbicides

Trade Name	Common Name	Rate of application/ha
Sofit	Pretilachlor 300 g/lit.EC	1.6 lit.
Goal	Oxyflorfen 240 g/lit.EC	0.5 lit.

##### Broad leaves/Sedge Herbicide

Trade Name	Common name	Rate of application/ha
Hedonal	2.4 D 550 g/lit.SL	0.95 - 1.2 lit.
	MCPA 600 g/lit.SL/EC	0.8 - 1.1lit.

Source: <http://www.doa.gov.lk/rrdi/index.php/en/rice>

	<p><b>New Rathna Rice Mill</b></p> <p><i>South Asia's Largest Rice Production Line</i></p> <p><b>Head Office &amp; Factory 01</b></p> <p>New Rathna Rice (PVT) Ltd. No. 54, Somawathiya Road, Pulasthigama, Polonnaruwa, Sri Lanka. Tel - 027-2242627 Email - newrathna.rice@gmail.com</p>
---	--

## The Rice Grain

The rice grain has both physical and chemical characteristics.

### Physical Structure

A rice grain is made up of an outside husk layer, a bran layer, and the endosperm, see Figure 1. The husk layer (lemma and palea) accounts for 20% of the weight of paddy and helps protect the grain kernel from insect and fungal attack. When the husk is removed, the rice is called brown rice. Brown rice contains the bran layer and the endosperm. The bran layer is made up of the pericarp and testa, the aluerone layer and the embryo. The degree to which this bran layer is removed is known as the milling degree. The desired amount of bran removed depends on the country. In Japan, the aluerone layer is often not removed however in many other countries all bran layers are removed to give very highly polished rice. The storage life of milled rice is improved when all of the bran layers are removed.

Physical characteristic	Percentage
Paddy	100
Husk	20
Brown rice	80
Meal Pericarp and testa (5-6%) Aluerone (1%) Embryo (3%)	8-10
White rice	70-72


### Chemical Composition of Milled Rice

Rice at 12% moisture contains approximately 80% starch and 7% protein. (Currey, 1984) Starch occurs in the endosperm as small many-sided granules while protein is present as particles that lie between the starch granules. Rice grain also contains sugars, fat, dietary fiber and minerals.

Component	Brown rice	White rice	Bran
Water (%)	13-14	13-14	13-14
Starch (%)	68-70	80	9
Amylose	28-30	33	6
Protein (%)	6-8	6-7	14
Fat	3	1	20
Fiber	2-3	0.5	25
Crude ash	1-1.5	0.5	9-10

Source: <http://www.newrathnarice.com/>

- Araliya Rice Mill**  
 3,2 ★★★★★ (4) · Industriegebiet  
 Polonnaruwa, Sri Lanka  
 +94 27 2 227227  
[WEBSITE](#) [ROUTE](#)  
 Jetzt geschlossen
- Chinthaka Rice Mill**  
 5,0 ★★★★★ (3) · Lebensmittelanbieter  
 Sri Lanka  
 +94 71 446 2349  
 Jetzt geschlossen
- Fazi Hajiyar Rice mill**  
 Keine Rezensionen · Reismühle  
 Sri Lanka  
[ROUTE](#)
- New Rathna Rice (pvt) Ltd.**  
 4,7 ★★★★★ (3) · Mühle  
 Polonnaruwa, Sri Lanka  
 +94 27 2 242627  
[WEBSITE](#) [ROUTE](#)  
 Jetzt geschlossen
- Sanjeewa Rice Mills**  
 4,3 ★★★★★ (3) · Lebensmittelanbieter  
 Malabe, Sri Lanka  
 +94 11 2 412703  
 Jetzt geschlossen
- Raveesha Rice Mill**  
 5,0 ★★★★★ (1) · Mühle  
 Sri Lanka  
 +94 71 566 0812  
[ROUTE](#)
- Ratna Rice Mill**  
 Keine Rezensionen · Mühle  
 Mawanella, Sri Lanka  
 +94 35 2 246217  
[ROUTE](#)



Source: <https://www.google.de>



ELSEVIER

available at [www.sciencedirect.com](http://www.sciencedirect.com)



journal homepage: [www.elsevier.com/locate/agwat](http://www.elsevier.com/locate/agwat)



## Predicting the impacts of climate change—A case study of paddy irrigation water requirements in Sri Lanka

C.S. De Silva <sup>a,\*</sup>, E.K. Weatherhead <sup>b</sup>, J.W. Knox <sup>b</sup>, J.A. Rodriguez-Diaz <sup>c</sup>

<sup>a</sup>Department of Agricultural Engineering, Faculty of Engineering Technology, The Open University of Sri Lanka, Nawala, Nugegoda, Sri Lanka

<sup>b</sup>Cranfield University, Bedford MK43 0AL, UK

<sup>c</sup>IFAPA, Junta de Andalucía, CIFA Alameda del Obispo, 14080 Cordoba, Spain

### ARTICLE INFO

#### Article history:

Received 3 March 2006

Accepted 13 June 2007

Published on line 23 July 2007

#### Keywords:

Food production

Geographical information system

Water resources

Spatial variability

Adaptation

### ABSTRACT

Nearly 72% of paddy production, the staple food in Sri Lanka, is grown during the wet season in dry areas where water resources are already stressed. Climate change datasets for Sri Lanka were derived using outputs from the UK Hadley Centre for Climate Prediction and Research Model (HadCM3) for selected scenarios for the 2050s, chosen from the Intergovernmental Panel of Climate Change Special Emission Scenarios Report. Water balance modelling and a geographical information system were used to model and map the impacts on irrigation requirements for wet season paddy. We examined two scenarios. The A2 scenario represents a heterogeneous, regionalised, market-led world, with high population growth, leading to a rapid increase in atmospheric carbon dioxide levels. The B2 scenario follows a similar regionalised future but with more moderate population growth and more concern for the environment and local sustainability, and a slower rate of increase in atmospheric carbon dioxide.


Results suggests that, during the wet season, average rainfall decreases by 17% (A2) and 9% (B2), with rains ending earlier, and potential evapotranspiration increasing by 3.5% (A2) and 3% (B2). Consequently, the average paddy irrigation water requirement increases by 23% (A2) and 13% (B2).

Mapping with GIS highlights the importance of considering spatial variation. Climate change impacts on wet season paddy production are positive in the extreme south, confirming results of a previous study. However, the impacts are negative across most of Sri Lanka. The adaptations needed are different in the two regions. Furthermore, spatial variation points to a further adaptation; the transfer of some paddy production to positively affected areas, which would not have been so clear if only point modelling had been used.

© 2007 Elsevier B.V. All rights reserved.

Source: <http://www.sciencedirect.com/science/article/pii/S037837740700162X>

Data for Tea production and processing



<https://upload.wikimedia.org/wikipedia/commons/1/13/Teaprocessing.svg>

[https://en.wikipedia.org/wiki/Tea\\_processing](https://en.wikipedia.org/wiki/Tea_processing)

[http://www.fao.org/fileadmin/templates/est/Climate\\_change/kenya/Navaisha\\_wkp\\_1/FAO\\_Uni\\_Manchester\\_AZAPAGIC\\_LCA\\_of\\_tea.pdf](http://www.fao.org/fileadmin/templates/est/Climate_change/kenya/Navaisha_wkp_1/FAO_Uni_Manchester_AZAPAGIC_LCA_of_tea.pdf)

## Dairy production

[http://sites.nationalacademies.org/cs/groups/pgasite/documents/webpage/pga\\_065995.pdf](http://sites.nationalacademies.org/cs/groups/pgasite/documents/webpage/pga_065995.pdf)

# U.S. Dairy Life Cycle Assessment

From Grass to Glass

Dr. Ying Wang  
National Academy of Sciences  
November 17, 2011



HEALTHY PEOPLE • HEALTHY PRODUCTS • HEALTHY PLANET

<http://www.sciencedirect.com/science/journal/09586946/31/supp/S1>

 **Export** ▾

 **Open Access articles** ▾

**Editorial Board**

Open Access 

*Page IFC*

 PDF (46 K)

### Editorial

**Carbon and water footprint of U.S. milk, from farm to table – Special issue: Editorial**

Open Access 

*Pages S1-S2*

Gregory D. Miller, Ying Wang

 PDF (89 K)

### Research Papers

**Greenhouse gas emissions from milk production and consumption in the United States: A cradle-to-grave life cycle assessment circa 2008** Original Research Article

Open Access 

*Pages S3-S14*

Greg Thoma, Jennie Popp, Darin Nutter, David Shonnard, Richard Ulrich, Marty Matlock, Dae Soo Kim, Zara Neideman, Nathan Kemper, Cashion East, Felix Adom

▶ [Abstract](#) |  PDF (799 K)

**Collecting complex comprehensive farm level data through a collaborative approach: A framework developed for a life cycle assessment of fluid milk production in the US** Original Research Article

Open Access 

*Pages S16-S20*

J.S. Popp, G.J. Thoma, J. Mulhern, A. Jaeger, L. LeFranc, N. Kemper

▶ [Abstract](#) |  PDF (546 K)

**Carbon footprint analysis of dairy feed from a mill in Michigan, USA** Original Research Article

Open Access 

*Pages S21-S28*

Felix Adom, Charles Workman, Greg Thoma, David Shonnard

▶ [Abstract](#) |  PDF (707 K) | [Supplementary content](#)

**Regional carbon footprint of greenhouse gas emissions from USA dairy farms: A cradle-to-farm gate assessment of the**

Open Access 

<https://nexus.openlca.org/search/query=milk>



# openLCA Nexus

Your source for LCA data sets.

Search

Options

## Database more

- GaBi 226
- ESU World Food 122
- Social Hotspots 113
- Agribalyse 72
- soca 63
- ecoinvent 63
- EuGeos' 15804-IA 56
- exiobase 48
- PSILCA 38
- ELCD 31
- more...

## Country more

- France 260
- Germany 247
- Romania 205
- Switzerland 200
- United Kingdom 196
- Netherlands 194
- Sweden 188
- Spain 188
- Slovenia 188
- Slovakia 188
- more...

766 data sets in 4 ms

### milk production, from cow (CA-QC)

Databases: ecoinvent v2.2 & v.3.3, soca v.1

System model: APOS, cut-off

Category: A:Agriculture, forestry and fishing/01:Crop and animal production, hunting and related service activities/0141:Raising of cattle and buffaloes

Version (internal): 03.03.000 Location: CA-QC

### yogurt production, from cow milk (CA-QC)

Databases: ecoinvent v2.2 & v.3.3, soca v.1

System model: APOS, consequential long-term, cut-off

Category: C:Manufacturing/10:Manufacture of food products/105:Manufacture of dairy products/1050:Manufacture of dairy products

Version (internal): 03.01.000 Location: CA-QC

### milk production, from cow (RoW)

Databases: ecoinvent v2.2 & v.3.3, soca v.1

System model: APOS

Category: A:Agriculture, forestry and fishing/01:Crop and animal production, hunting and related service activities/0141:Raising of cattle and buffaloes

Version (internal): 03.02.000 Location: RoW

### yogurt production, from cow milk (CA-QC)

<http://www.sciencedirect.com/science/article/pii/S221431731500030X>



Download PDF

Export



## Information Processing in Agriculture

Volume 2, Issue 2, September 2015, Pages 101-108

open access



### Life Cycle Assessment modeling of milk production in Iran

Hamzeh Soltanali<sup>a</sup>, Bagher Emadi<sup>✉, a</sup>, Abbas Rohani<sup>a</sup>, Mehdi Khojastehpour<sup>a</sup>, Amin Nikkhah<sup>✉, a, b</sup>

**Show more**

<https://doi.org/10.1016/j.inpa.2015.06.003>

[Get rights and content](#)

Open Access funded by China Agricultural University

Under a Creative Commons [license](#)

Highlights